

Gene conferring resistance to the antibacterial 4,5-dihydroxy-2-cyclopenten-1-one (DHCP),
the protein encoded by same, and applications thereof

Sub
A107

JC918 U.S. PTO
09/805681
03/14/01

Inventors:

Sangita Phadtare
Kunitoshi Yamanaka
Ikunoshin Kato
Masayori Inouye

SCANNED, # 10

Sub A10 > **Gene conferring resistance to the antibacterial 4,5-dihydroxy-2-cyclopenten-1-one (DHCP), the protein encoded by same, and applications thereof**

This patent application claims the benefit of U.S. Provisional Application No. 60/228,727, filed 8/29/2000. This earlier provisional application is hereby incorporated by reference.

In spite of a number of antibiotics available against a variety of bacteria, due to emergence of multiple drug resistant strains, the search for newer and more effective antibacterial compounds has continued. 4,5-dihydroxy-2-cyclopenten-1-one (DHCP) (see Fig.1) is a compound having antibacterial activity against a variety of gram-positive and -negative bacteria, such as *Escherichia coli*, *Bacillus*, *Salmonella*, *Staphylococcus* etc. The process for manufacture and the properties of DHCP have been patented (Koyama *et al.*, 1999). It is prepared by the heat-treatment of uronic acid or its derivatives, wherein uronic acid is galacturonic acid, glucuronic acid, mannuronic acid or iduronic acid. It is also produced from roasted or parched vegetables, fruits, cereals, mushrooms, sea algae, cortex or cartilage. It has been demonstrated that this compound induces cancer cell differentiation and apoptosis. It has potential application as therapeutic or preventive agent against cancer and also as an antibacterial agent in antiseptics, dentrifices, cosmetics and bathing agents (Koyama *et al.*, 1999).

We have isolated a multicopy suppressor from an *E. coli* genomic library for the DHCP toxicity. The putative protein encoded by this gene showed high homology to known efflux proteins conferring resistance to a number of antibiotics including chloramphenicol, bicyclomycin and tetracycline. The gene was mapped at 37.5 min on the *E. coli* chromosome. It is designated as *dep* for DHCP efflux protein. However, the Dep protein does not confer cross-resistance to any of the antibiotics tested.

10/23/12
KR


~~SUMMARY~~
~~BRIEF DESCRIPTION OF THE INVENTION~~

The provisional application describes the cloning of a gene encoding a transmembrane protein from *E. coli*. This protein, when expressed from a multi-copy plasmid, functions to transport 4,5-dihydroxy-2-cyclopenten-1-one (DHCP) out of the cell.


5 DHCP and functionally equivalent compounds are represented by the formulas [I] and [II] and include optically active compounds thereof. In Formula I, R^1 and R^2 are the same or different and each of them is hydrogen, a straight or branched alkyl group, a straight or branched alkenyl group, an aromatic group, an aromatic-aliphatic group, with the proviso that R^1 = a benzyl group and $R^2 = H$ is excluded.

See References 3 and 4. R^3 - R^6 are independently hydrogen or an alkyl group, preferably a lower alkyl group such as a C_1 - C_6 alkyl.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
21



Formula I



Formula II

In Formula II, R^1 and R^2 are the same or different and each of them is hydrogen, a straight or branched alkyl group, a straight or branched alkenyl group, an aromatic group, an aromatic-aliphatic group, with the proviso that the case where $R^1 = R^2 = \text{CH}_3$ is excluded. See References 5 and 6. R^3 - R^6 are independently hydrogen or an alkyl group, preferably a lower alkyl group such as a C_1 - C_6 alkyl.

DHCP is shown to possess anti-bacterial activity; it inhibits cell growth at a concentration of $350\mu\text{M}$ or higher. At lower concentrations, it causes cells to elongate and grow poorly. To determine if *E. coli* is naturally resistant to DHCP, a library of *E. coli* genomic DNA fragments was transformed into strain JM83 and grown on agar plates containing $400\mu\text{M}$ DHCP. Colonies that were capable of growing on this medium were isolated. DNA was isolated from these colonies to identify and sequence the cloned genomic fragment that specified resistance. Four genes were found in the fragment that conferred resistance. Inactivation of various combinations of these four genes led to the conclusion that ORF389 was responsible for conferring resistance. This was confirmed by cloning ORF389 by itself into pUC19 (a multi-copy plasmid) and transforming strain JM83. The resultant cells were resistant to DHCP.

Comparison of the nucleotide sequence of ORF389 with the *E. coli* gene database showed that it was similar to known efflux proteins involved in conferring resistance to chloramphenicol and other antibiotics. Further analysis of the predicted structure of the protein encoded by ORF389 suggested that it was a membrane protein; it possesses multiple transmembrane domains and shares structural similarity with the aforementioned chloramphenicol efflux polypeptides.

To determine if ORF389 was capable of conferring resistance to other antibiotics such as chloramphenicol, spectinomycin, and tetracycline, the transformed JM83 cells containing the pUC19/ORF389 plasmid were plated on media containing these antibiotics. The presence of ORF389 failed to confer resistance to any antibiotic other than DHCP, suggesting that the efflux activity of the Dep protein is specific for DHCP.

It is important to note that ORF389 confers resistance to DHCP only when it is present in multiple copies in the cell. The gene is naturally found in the genome of *E. coli* cells, but it is present in single copy. Such cells are susceptible to the antimicrobial activity of DHCP. When ORF389 is cloned into pUC19 and introduced into JM83 cells, it is present in multiple copies (up to several hundred copies of the gene per cell), since pUC19 is maintained in up to several hundred copies per cell. Only when the gene dosage is increased, is resistance to DHCP found. The mechanism of resistance is simply increased efflux activity arising from the increased expression of the efflux protein in the transformed cells.

It should be noted that due to the degeneracy of the genetic code, the nucleotide sequence encoding an efflux protein that is responsible for conferring resistance to DHCP or a compound functionally equivalent to DHCP may vary from the nucleic acid sequence disclosed herein.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is the chemical structure of 4,5-dihydroxy-2-cyclopenten-1-one (DHCP).

FIG. 2A is a graphical representation of the effect of DHCP concentration on the growth
5 of *E. coli*.

FIG. 2B is a graphical representation of the effect of DHCP concentration on the survival
of *E. coli*.

FIG. 3 is a restriction mapping of the plasmid pSP001 showing the DNA fragments
conferring resistance to DHCP.

FIG. 4 is a comparison of the amino acid sequence of the polypeptide encoded by *dep*
with the proteins encoded by *cmr*, *cmrA*, *cmx*, *cmlv*, *bcr*, *bmr3*, *yjcC*, and *tet*.

FIG. 5 is a comparison of the hydropathic profiles of the putative proteins encoded by
dep, *cmr*, and *cml*.

FIG. 6 is a nucleotide sequence showing the DNA sequence of a region of the *E. coli*
genome containing the sequence of the *dep* gene.

FIG. 7 is a nucleotide sequence showing the isolated DNA sequence of the *dep* gene.

10/23/02
KK

DETAILED
~~ADDITIONAL~~ DESCRIPTION OF THE INVENTION

Mode of Action

DHCP is a compound that exhibits antimicrobial and anti-tumor activity. It is made by heating various uronic acids (e.g., glucuronic acid, galacturonic acid, mannuronic acid). The inventors have cloned a gene from *E. coli* that encodes a protein which is capable of transporting DHCP out of the cell. This transport protein shows sequence similarity with known efflux proteins that function to transport antibiotics such as chloramphenicol out of the cell. It has been shown that organisms which overexpress the transport protein become resistant to DHCP, probably because they are able to efficiently transport DHCP. Overexpression of the transport protein arises from the presence of multiple copies of the gene, rather than increased expression from the endogenous gene in *E. coli*. In other words, all *E. coli* possess a single copy of the transport gene. However, the level of transport protein expression from a single copy of the gene is insufficient to confer resistance to DHCP. The inventors have cloned the gene into a high copy number plasmid, pUC19, which is maintained in *E. coli* cells at 200-500 copies per cell. Thus, transformed *E. coli* containing this plasmid construct will possess 200-500 copies of the transport gene, and protein expression from multiple copies is greater than from a single copy. These transformed cells are resistant to DHCP.

The general mode of action of DHCP requires that it enter the target cell. Resistance to DHCP can occur if DHCP is transported out of the cell as fast as or faster than it enters the cell. Given that, the concentration of DHCP within the cell can never accumulate to a toxic dose and the cell is resistant to the antimicrobial effects of the compound. Apparently, the transport protein encoded by gene disclosed does not transport DHCP very efficiently, or the amount of transport protein expressed from the endogenous gene is very low. In either case, the presence of more transport protein (arising from many copies of the gene) will result in more efficient transfer of DHCP out of the cell.

Applications

An important application of the gene of the invention will be its use in studies to identify inhibitors of efflux activity. Such inhibitory compounds will function to block the transport activity. Thus a microbe or a tumor cell that is resistant to DHCP can be made to be more sensitive to the compound by preventing the resistant cell from transporting the compound back out. It is also conceivable that inhibitors of the transport gene of the invention may also be active in blocking transport of other efflux proteins such as the efflux proteins that transport chloramphenicol, or the P glycoprotein family of multiple drug resistant proteins. The P glycoproteins are expressed in many tumor cells, making these tumors resistant to chemotherapy agents. Abstracts regarding studies of P glycoproteins are referenced ~~above~~ ^{herein}.

KK
10/26/42

KK
10/23/02

~~DETAILED DESCRIPTION OF THE DRAWINGS~~

FIG. 1 is the chemical structure of 4,5-dihydroxy-2-cyclopenten-1-one (DHCP).

FIG. 2 Effect of DHCP concentrations on the growth of *E. coli*.

5 A. The JM83 cells were grown in LB medium up to Klett unit of 50 and DHCP was added at various concentrations (0-400 μ M). After the growth reached to Klett unit of 90-100, cells were diluted with medium containing respective concentrations of DHCP and growth was further monitored. DHCP concentration: 0 μ M, open squares; 50 μ M, closed diamonds; 100 μ M, open circles; 250 μ M, open diamonds; 400 μ M closed squares.

10 B. Overnight grown cells of *E. coli* JM83 were diluted appropriately and plated on LB plates containing different concentrations of DHCP (0-350 μ M). The number of colonies on the plate without DHCP was taken as 100% and the other numbers were expressed as relative percentages.

15 FIG. 3 Restriction mapping of the plasmid pSP001 conferring resistance to DHCP. Four ORFs comprising the DNA fragment (5.2 kb) conferring resistance to DHCP and the flanking ORFs are shown. The orientation of each ORF is marked with an arrow. The restriction enzyme sites are also shown. The ORFs are not drawn to scale. The plasmid pSP001 containing the DNA fragment conferring resistance to DHCP was digested with restriction enzymes to disrupt each of four ORFs, religated and transformed into JM83 cells. The transformants were then examined for their sensitivity to DHCP (400 μ M). The enzymes used
20 for digestion were: for *purR*: *Mlu*I for *ydhB*; *Nru*I-*Eco*47III, for ORF389, *purR*, and *ydhB*: *Nru*I and *Sma*I, for ORF389: *Ava*I and for *purR* and *ydhB*: *Mlu*I and *Nru*I. For construction of plasmid with ORF389 (*dep*), the plasmid pSP001 was digested with *Sma*I and *Msc*I, the fragment was purified and cloned into pUC19 to yield plasmid pSP007.

25 FIG. 4 The sequence homology between Dep, Cmr, CmrA, Cmx, CmlV, BcR, Bmr3, YjcC and Tet. Identical and similar sequences are marked with black and gray boxes, respectively. The consensus sequences for transmembrane proteins are marked with dotted lines and are represented as I, II, and III stretches.

Sgt

FIG. 5 Hydropathic profiles of Dep (A), Cmr from *Rhodococcus faciens* (B) (6) and Cml from *Streptomyces lividans* (C) (8). Horizontal bars indicate predicted transmembrane regions.

Sub A2
5
FIG. 6 is a nucleotide sequence showing the DNA sequence of a region of the *E. coli* genome containing the sequence of the *dep* gene. This region of the *E. coli* genome is available at Accession No. AE000261 U00096. The sequence shown is that of nucleotides 4381-8280. The *dep* gene is encoded by nucleotides 4627-5838. The *dep* sequence is shown in brackets.

Sub A3
10
FIG. 7 is a nucleotide sequence showing the isolated DNA sequence of the *dep* gene. The plasmid pSP007 was confirmed to contain the *dep* gene by obtaining DNA sequence data from one end of the 1.7 kb insert. Sequence data obtained in this manner matched the first.

~~Detailed Description of Experimental Work~~

Effect of DHCP on the growth of *E. coli*

KK
10/23/02
10
15
20
25
The *E. coli* wild-type strain JM83 [F^- *ara* Δ (*lac-proAB*) *rpsL*(*str'*)] (Yanisch-Perron *et al.*, 1985) was grown in Luria broth (LB). Media were supplemented with ampicillin (final concentration of 50 μ g/ml) whenever required. To check the effect of DHCP on the growth of *E. coli*, cells grown overnight in LB medium were diluted into fresh LB medium. After the growth reached to the Klett unit of 50, DHCP was added at various concentrations (0-400 μ M) and growth was further monitored. After it reached to the Klett unit of 90-100, it was diluted 10-fold into media containing respective concentrations of DHCP. Fig. 2A shows the effect of different concentrations of DHCP on *E. coli*. The growth was slowed after 3 h of incubation in the presence of 50 μ M DHCP, but it reached the maximum density after 8 h, similar to that without DHCP. The cells grew more slowly after 3 h incubation with 100 μ M of DHCP and the maximum cell density was lower than that without DHCP. In the presence of 250 μ M DHCP, growth was severely impaired after 3 h of incubation and cells stopped growing after 5 h. In the presence of 400 μ M DHCP, cell growth stopped after 4 h of incubation. Microscopic examination of the cells grown with 250 μ M DHCP for 8 h showed that the cells were elongated

forming filaments, which are approximately 15-fold longer than the control cells. DAPI (diamidino phenylindole)(Hiraga *et al.*, 1989) staining of these cells showed that the chromosomal condensation of the cells might be impaired by DHCP (data not shown).

To check the colony formation ability of *E. coli* at various concentrations of DHCP, cells grown overnight in LB medium were diluted appropriately and plated on LB plates containing DHCP (0-350 μ M). After incubation at 37 $^{\circ}$ C, the number of colonies on each plate were counted. The number of colonies on the control plate without DHCP was taken as 100% and the other numbers were expressed as relative percentages (Fig. 2B). In the presence of 300 μ M DHCP, 100-fold decrease in the colony numbers was observed. When 1×10^4 cells were plated on LB medium containing 350 μ M DHCP, no colonies were obtained.

Screening of an *E. coli* genomic library for genes conferring resistance to DHCP

In order to examine if *E. coli* contains a gene(s) that confers resistance to DHCP, the *E. coli* genomic library was screened. The construction of *E. coli* genomic library was described previously (Lu and Inouye, 1998). The partially digested *Sau*3AI chromosomal DNA fragments from *E. coli* JM83 were cloned into the *Bam*HI site of pUC19. The JM83 cells were transformed with the genomic library. Transformants were isolated for their ability to grow on DHCP (400 μ M) containing LB plates at 37 $^{\circ}$ C. Plasmid DNA was isolated from the resistant colonies, purified and retransformed into JM83 cells to confirm its ability to confer resistance to DHCP. The plasmid was designated as pSP001 and was found to contain a 5.2-kb DNA fragment. This fragment was sequenced from both ends using Sequenase and BLAST search was carried out for the analysis of homology of this fragment with the entire *E. coli* genome. It was found that this DNA fragment is located at 37.5 min on the *E. coli* chromosome and contains four ORFs (Fig. 3): ORF389, *purR* encoding purine synthesis repressor, *ydhB* encoding a homologue of the *cyn* operon transcriptional activator and *ydhC* encoding a homologue of bicyclomycin resistance protein (Berlyn *et al.*, 1996).

To determine which gene is responsible for conferring resistance to DHCP, several deletion constructs were prepared as shown in Fig. 3. Disruption of *purR*, *ydhB* and both *purR*

and *ydhB* had no effect on the resistance to DHCP (constructs pSP002, pSP003 and pSP006, respectively). However, disruption of ORF389 with *purR* and *ydhB* (pSP004) as well as disruption of ORF389 alone (pSP005) resulted in loss of DHCP resistance. We thus cloned ORF389 separately in pUC19 (pSP007), transformed the resultant plasmid in JM83 and checked sensitivity to DHCP. This plasmid conferred resistance to DHCP. These results clearly demonstrate that ORF389 is responsible for resistance to DHCP when cloned in a multicopy plasmid and further work was carried out using the plasmid pSP007. The ORF389 was named as *dep* – DHCP efflux protein (see below).

Homology analysis of ORF389 with other genes conferring drug resistance

Using BLAST-homology search computer program, we carried out a homology search for the putative protein encoded by *dep*. Fig. 4 shows nine proteins showing significantly high homology with *Dep*. Half of these proteins confer resistance to chloramphenicol. The proteins showing the highest degree of homology include: Cmr from *Rhodococcus fasciens* (Desomer *et al.*, 1992), CmrA from *R. erythropolis* (Nagy *et al.*, 1997), Cml from *Streptomyces lividans* 1326 (Dittrich *et al.*, 1991), Cmx from *Corynebacterium striatum* (Accession no. U72639), and CmlV from *S. venezuelae* ISP5230 (Mosher *et al.*, 1995). As seen from Fig. 4, *Dep* has the highest degree of homology with Cmr, product of chloramphenicol resistant gene (*cmr*) as compared to other proteins. Cmr protein was shown to contain three consensus sequences defined by Rouch *et al.* (1990) for transmembrane proteins. These sequences are at similar positions with respect to the predicted transmembrane domains. These are marked in Fig. 6 with dotted lines and are designated as I, II, III. In case of *Dep*, the first stretch (I) comprising of LP is completely homologous with the stretch defined by these authors. The second stretch (II) shows 50% similarity with that of Cmr protein and the third stretch (III) is homologous between these two proteins except for one residue. According to the model proposed by Rouch *et al.* (1990), the stretches I and III are located on the outside of the cytoplasmic membrane and the stretch II is located on the inside of the membrane. The positions of the membrane loops for the putative

protein encoded by *qacA* were ascertained by inspecting the antigenic index profile and turn prediction. Such regions have a high antigenic index and turn probability (Rouch *et al.*, 1990).

In addition to homology in the primary sequences, the hydropathic profile of Dep (Fig. 5A) is significantly similar to those of Cmr of *R. faciens* (Desomer *et al.*, 1992) (Fig. 5B) and Cml of *S. lividans* (Dittrich *et al.*, 1991) (Fig. 5C). Dep is predominantly hydrophobic and probably contains 12 predicted transmembrane α -helices (Fig. 5A).

The other proteins homologous to Dep include BcR (bicyclomycin- resistance protein) from *E. coli* (Bentley *et al.*, 1993), Bmr3 from *B. subtilis* involved in the multiple drug efflux pump conferring resistance to puromycin, tosylflouxacin, norfloxacin (Ohki and Murata, 1997), Tet from *Staphylococcus hyicus* conferring tetracycline resistance (Schwarz *et al.*, 1992) and YjcC conferring tetracenomycin-resistance (Accession no. D90826) (Fig. 4). All of these are efflux proteins, which is one of the most common mechanisms for drug resistance. We speculate that *dep* encodes a putative efflux protein that forms a cytoplasmic channel specific for DHCP. The homologies are more prominent towards the N-terminal end of the proteins, which also is a common feature for efflux proteins (Desomer *et al.*, 1992).

Measurement of minimum inhibitory concentrations for cells harboring pUC19 and pSP007

Since Dep shows homology to efflux proteins for multiple drug resistance, we checked if it confers resistance to other antibiotics as well. The *E. coli* wild-type cells harboring pUC19 or pSP007 plasmid were grown overnight in LB medium containing ampicillin. The cells were diluted 10- and 1000- times, and 5 μ l of each dilution (corresponding to 3.5×10^5 cells and 3.5×10^3 cells, respectively) was spotted on LB plates containing serial dilutions of kanamycin, chloramphenicol, spectinomycin, tetracycline and DHCP. Plates were incubated at 37 $^{\circ}$ C for 20 h. As seen from Table1, pSP007 did not confer significant cross-resistance to any of the antibiotics tested. The MIC values for cells harboring pUC19 and pSP007 were same for spectinomycin, chloramphenicol and tetracycline. The MIC value was two times higher for kanamycin for the cells harboring pSP007 than the cells with pUC19. The MIC value for DHCP

on the other hand was 8 times higher for the cells harboring pSP007 than that for the cells with pUC19. It is interesting that Dep did not confer resistance to chloramphenicol, in spite of the high homology to *cmr*.

Table 1. Minimum inhibitory concentrations (MICs) of various antibiotics for *E. coli* JM83 cells harboring pUC19 and pSP007.

	MICs (μg/ml)				
	kanamycin	spectinomycin	chloramphenicol	tetracycline	DHCP
cells with pUC19	25	12.5	6.25	3.125	25
cells with pSP007	50	12.5	6.25	3.125	200

MICs for both dilutions of the cells (3.5×10^5 and 3.5×10^3 cells) were the same.

Prior Art References

United States Patent No. 6,087,401 to Koyama, et al. Cyclopentones, process for preparing same, and the use thereof.

5

This patent discloses a method of manufacturing 4,5-dihydroxy-2-cyclopenten-1-one (DHCP). It also describes the antibacterial activity of DHCP.

In contrast, the invention disclosed in the present provisional application relates to a gene, *dep*, that, when present in multiple copies in bacterial cells, confers resistance to the antibacterial activity of DHCP, thus rendering the bacteria resistant to killing by DHCP. The present application also describes the protein encoded by the *dep* gene.

European Patent EP 0 941 981 A1 to Koyama, et al. Cyclopentones, process for preparing same, and the use thereof.

This patent application relates to essentially the same subject matter as that described in United States Patent No. 6,087,401 to Koyama, et al.

United States Patent No. 6,111,145 to Kobayashi et al. Cyclopentenone derivative.

This patent relates to functionally equivalent ether derivatives of DHCP and discloses the biological activity of these derivatives.

European Patent Publication EP 1 000 923 A1 to Kobayashi et al. Cyclopentenone derivatives.

This patent application relates to essentially the same subject matter as that described in United States Patent No. 6,111,145 to Kobayashi et al.

United States Patent No. 6,136,854 to Koyama et al. Cyclopentenone derivative.

5

This patent relates to functionally equivalent ester derivatives of DHCP and discloses the biological activity of these derivatives.

European Patent Publication EP 0 976 717 A1 to Koyama et al. Cyclopentenone derivatives.

This patent application relates to essentially the same subject matter as that described in United States Patent No. 6,136,854 to Koyama et al.

Clinical significance of P-glycoprotein expression and function for response to induction chemotherapy, relapse rate and overall survival in acute leukemia. C. Wuchter, et al. Haematologica 85(7):711-21 (2000).

In acute leukemia, a multidrug-resistance (MDR) phenotype mediated by P-glycoprotein (P-gp) contributes to chemotherapy failure. This study investigated whether P-gp expression levels or functional P-gp activity was a better predictor of response to induction chemotherapy, relapse rate and overall survival in acute leukemia. The data demonstrated that the functional rhodamine-123- (rh123)-efflux assay was preferred over P-gp expression analysis by monoclonal antibodies in acute leukemia.

Increased drug delivery to the brain by P-glycoprotein inhibition. A.J. Sadeque, et al. Clinical Pharmacology & Therapeutics 68(3):231-7 (2000).

In vitro studies had demonstrated that the antidiarrheal drug loperamide is a substrate for the efflux membrane transporter P-glycoprotein. Although loperamide is a potent opiate drug, it does not opiod central nervous system effects, such as respiratory depression, when given to patients at usual doses. This study tested the hypothesis that inhibition of P-glycoprotein with quinidine would increase the entry of loperamide into the central nervous system, thus causing respiratory depression. The results demonstrated that although loperamide produced no respiratory depression when used alone, respiratory depression was seen when loperamide was administered with quinidine.

Expression of the multidrug-resistance-associated protein in myelodysplastic syndromes. S. Poulain, et al. British Journal of Haematology 110(3):591-8 (2000).

In myelodysplastic syndromes (MDS), P-glycoprotein (P-gp) expression is associated with drug resistance, while the clinical significance of the multidrug resistance-associated protein (MRP1) is unclear. In this study of bone marrow from patients with MDS, expression of MRP1 was correlated with disease stage in MDS. With respect to P-gp, discordant expression/function of MRP1 was found in some cases, suggesting the existence of nonfunctional transport proteins in MDS. MRP1 expression did not appear to be a prognostic factor in MDS.

Soft tissue leiomyosarcomas and malignant gastrointestinal stromal tumors: differences in clinical outcome and expression of multidrug resistance proteins. B. E. Plaat, et al. Journal of Clinical Oncology 18(18):3211-20 (2000).

In this study, parameters associated with multidrug resistance (MDR) were compared between soft tissue leiomyosarcomas (LMS) and malignant gastrointestinal stromal tumors (GIST). Immunohistochemistry was used to detect P-glycoprotein (P-gp), multidrug resistance protein (MRP(1)), lung resistance protein (LRP), and c-kit. The results demonstrate that LMS

patients have better survival rates compared to GIST patients, and the pattern of metastasis differs between the two patient groups. The expression of the MDR proteins tested is less pronounced in LMS than in GIST.

5 Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: A new family of genes responsible for autoinducer production. M.G. Surette, et al. Proc. Natl. Acad. Sci. 96:1639-44 (1999).

In bacteria, the regulation of gene expression in response to changes in cell density, called quorum sensing, is dependent on hormone-like molecules known as autoinducers that are produced by the bacteria and accumulate in the external environment as the bacterial cell population increases. The marine bacterium *Vibrio harveyi* has been shown to have two parallel quorum sensing systems, each composed of a sensor-autoinducer pair. The two different autoinducers belonging to each system have been termed autoinducer 1 (AI-1) and autoinducer 2 (AI-2). The identification and analysis of the genes responsible for AI-2 production in *E. coli*, *S. typhimurium*, and *V. harveyi* is reported.

Quorum sensing in *Vibrio fischeri*: Probing autoinducer-LuxR interactions with autoinducer analogs. A. L. Schaefer, et al. Journal of Bacteriology 178:2897-2901 (1996).

20 In *Vibrio fischeri*, luminescence genes are activated by the transcription factor LuxR in combination with a diffusible signal compound known as the autoinducer. This study analyzed the ability of a number of autoinducer analogs to interact with LuxR.

Regulation of quorum sensing in *Vibrio harveyi* by LuxO and Sigma-54. B.N. Lilley and B. L. Bassler. Molecular Microbiology 36(4):940-954 (2000).

The bioluminescent marine bacterium *Vibrio harveyi* controls light production (*lux*) by a quorum-sensing circuit. This study demonstrates that the response regulator protein LuxO functions as an activator protein via interaction with the alternative sigma factor, σ^{54} . Since LuxO is responsible for repression of the luciferase structural operon (*luxCDABEGH*), these results suggest that LuxO, together with σ^{54} , functions to activate a negative regulator of luminescence.

14. Bentley, J., Hyatt, L.S., Ainley, K., Parish, J.H., Herbert, R.B., and White, G.R. 1993. Cloning and sequence analysis of an *Escherichia coli* gene conferring bicyclomycin resistance. *Gene* 127: 117-120.
15. Berlyn, M.K.B., Low, K.B., and Rudd, K.E. 1996. Linkage map of *Escherichia coli* K-12, Ed. 9. Pages 1715-1902. In Neidhardt, F.C., Curtiss III, R., Ingraham, J.L., Lin, E.C.C., et al. (ed) *Escherichia coli and Salmonella. Cellular and Molecular Biology*, Vol. 2, 2nd Ed., ASM Press, Washington DC.
16. Desomer, J., Vereecke, D., Crespi, M., and Van Montagu, M. 1992. The plasmid-encoded chloramphenicol-resistance protein of *Rhodococcus fascians* is homologous to the transmembrane tetracycline efflux proteins. *Mol. Microbiol.* 6: 2377-2385.
17. Dittrich, W., Betzler, M., and Schrempf, H. 1991. An amplifiable and deletable chloramphenicol-resistance determinant of *Streptomyces lividans* 1326 encodes a putative transmembrane protein. *Mol. Microbiol.* 5:2789-2797.
18. Hiraga S., Niki, H., Ogura, T., Ichinose, C., Mori, H., Ezaki, B., and Jaffe, A. 1989. Chromosome partitioning in *Escherichia coli*: novel mutants producing anucleate cells. *J. Bacteriol.* 171:1496-505.
19. Koyama, N., Sagawa, H., Kobayashi, E., Enoki, T., Wu, H-K., Nishiyama, E., Ikai, K., and Kato, I. 1999. Cyclopentanones, process for preparing the same, and the use thereof. European Patent (EP 0941 981 A1, date of publication: 09/15/1999).

- KK
10/23/02
20. Lu Q., and Inouye, M. 1998. The gene for 16S rRNA methyltransferase (*ksgA*) functions as a multicopy suppressor for a cold-sensitive mutant of Era, an essential RAS-like GTP-binding protein in *Escherichia coli*. J. Bacteriol. 180:5243-5246.
21. Mosher, R.H., Camp, D.J., Yang, K., Brown, M.P., Shawl, W.V., and Vining, L.C. 1995. Inactivation of chloramphenicol by O-phosphorylation. J. Biol. Chem. 270:27000-27006.
22. Nagy, I., Schoofs, G., Vanderleyden, J., and De Mot, R. 1997. Transposition of the IS21-related element IS1415 in *Rhodococcus erythropolis*. J. Bacteriol. 179:4635-4638.
23. Ohki, R., and Murata, M. 1997. *bmr3*, a third multidrug transporter gene of *Bacillus subtilis*. J. Bacteriol. 179:1423-1427.
24. Rouch, D.A., Cram, D.S., DiBerardino, D., Littlejohn, T.G., and Skurray, R.A. 1990. Efflux-mediated antiseptic resistance gene *qacA* from *Staphylococcus aureus*: common ancestry with tetracycline- and sugar-transport proteins. Mol. Microbiol. 4:2051-2062.
25. Schwarz, S., Cardoso, M., and Wegener, H.C. 1992. Nucleotide sequence and phylogeny of the *tet(L)* tetracycline resistance determinant encoded by plasmid pSTE1 from *Staphylococcus hyicus*. Antimicrob. Agents Chemother. 36:580-588.
26. Yanisch-Perron, C., Vieira, J., and Messing, J. 1985. Improved M13 phage cloning vectors and host strains: Nucleotide sequences of the M13mp18 and pUC19 vectors. Gene 33: 103-119.

All references cited herein are incorporated by reference.